

Sharing clinical notes, and placebo and nocebo effects: Can documentation affect patient health?

Journal of Health Psychology
2022, Vol. 27(1) 135–146
© The Author(s) 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1359105320948588
journals.sagepub.com/home/hpq


Charlotte R Blease¹ , Tom Delbanco^{1,2},
John Torous^{1,2}, Moa Ponten³, Catherine M DesRoches^{1,2},
Maria Haglund^{1,4}, Jan Walker^{1,2‡} and Irving Kirsch^{1,2‡}

Abstract

This paper connects findings from the field of placebo studies with research into patients' interactions with their clinician's visit notes, housed in their electronic health records. We propose specific hypotheses about how features of clinicians' written notes might trigger mechanisms of placebo and nocebo effects to elicit positive or adverse health effects among patients. Bridging placebo studies with (a) survey data assaying patient and clinician experiences with portals and (b) randomized controlled trials provides preliminary support for our hypotheses. We conclude with actionable proposals for testing our understanding of the health effects of access to visit notes.

Keywords

placebo, placebo effects, electronic health records, clinical documentation, patient portals, nocebo effects

Introduction

Clinicians in more than a dozen countries currently offer patients online access to part, or all of the notes they write and house on computerized devices (Essén et al., 2018). The practice—known as “open notes”—is growing. In the United States, currently, over 50 million patients are offered access to such personal medical information via secure online portals, and in March 2020 the federal government released a new ruling stipulating that digital accessibility to patients' records will become mandatory. From April 2020, general practitioners working for NHS England will be obliged, on a prospective basis, to share the clinical notes that they write with their patients. (Richards, 2020) In

Sweden, where most patients can already read their clinical notes, all tax-funded health facilities were required to provide fully transparent patient access to electronic health records by 2020.

¹Beth Israel Deaconess Medical Center, Boston, MA, USA

²Harvard Medical School, Boston, MA, USA

³Karolinska Institute, Stockholm, Sweden

⁴Uppsala University, Uppsala, Sweden

‡Joint senior authors

Corresponding author:

Charlotte R Blease, OpenNotes, Division of General Medicine, Beth Israel Deaconess Medical Center, 133 Brookline Avenue, Boston, MA 02215, USA.
Email: cblease@bidmc.harvard.edu

In addition to legal, technical, and ethical challenges that pose ongoing obstacles to open notes (Blease et al., 2020a; 2020b; Hägglund et al., 2019; Mehta et al., 2019), many clinicians remain skeptical about the benefits to patients (Dobscha et al., 2016; Petersson and Erlingsdóttir, 2018b). Survey research reveals that many physicians fear access might cause confusion and anxiety among patients and could also disrupt workflow (Delbanco et al., 2012; Miller Jr et al., 2016; Petersson and Erlingsdóttir, 2018b). Mental health clinicians cite potential harms to psychiatric and psychotherapy patients if they were to read their clinical notes (Denneson et al., 2017; Dobscha et al., 2016; Petersson and Erlingsdóttir, 2018a). Yet, overall the majority of clinicians, including those who work primarily with mental illness, opine that sharing clinical notes with patients is a good idea (Delbanco et al., 2012; DesRoches et al., 2020; Petersson and Erlingsdóttir, 2018b).

In this paper we argue that both perspectives may be correct. Drawing on current findings in placebo studies and research into the practice of open notes, we hypothesize that, depending on their content, patient portal access to notes may generate both genuinely beneficial *and* genuinely adverse health effects by engaging perceptual and cognitive processes that give rise to placebo and nocebo effects, respectively. Specifically, we propose that the content and tone of clinical notes may influence expectancies via cues of clinician competence and empathy, and by documenting treatment rationale. Conditioning may also play a role.

We begin by providing a brief overview of research into these psychobiological pathways that are now recognized to elicit placebo and nocebo effects. Next, we propose four hypotheses describing how sharing clinical notes via patient portals might activate these mechanisms to modulate health effects. We then connect these hypotheses with current research into open notes and provide some evidence for the relationship between clinical note sharing, and placebo and nocebo effects. Finally, we suggest novel research designs and methodologies to expand this inquiry.

Mechanisms and pathways of placebo and nocebo effects

Empirical and conceptual inquiry into the role of placebo and nocebo effects has burgeoned (Colagiuri et al., 2015; Evers et al., 2018; Wolters et al., 2019), and in recent years a mature scientific research program in “placebo studies” has emerged (Blease, 2018; Blease and Annoni, 2019). A wealth of research shows that placebo effects are genuine psychobiological events that trigger top-down cognitive processes to elicit measurable physiological effects that can be clinically beneficial. Although investigators have examined only a limited number of symptoms and conditions, several are apparently susceptible to placebo effects (Kaptchuk and Miller, 2015). These include some of the most prevalent complaints and conditions for patients seeking primary care: depression and anxiety (Kirsch, 2019; Sugarman et al., 2014), pain (Amanzio et al., 2001; Locher et al., 2017), alcohol dependence (Weiss et al., 2008), and irritable bowel syndrome (Kaptchuk et al., 2008; Vase et al., 2005). Studies demonstrate that placebo effects can substantially augment the potency of active pharmacological treatments (Amanzio et al., 2001), and a recent consensus paper composed by experts in placebo studies proposed that harnessing the power of placebo effects for prevalent conditions is a medically worthwhile pursuit (Evers et al., 2018).

Research has predominantly focused on “response expectancies” as the core mechanism of placebo effects (Colagiuri et al., 2015; Jensen et al., 2012; Kirsch, 1985, 1997, 2018). The majority of studies investigating placebo effects have concentrated on patients’ consciously-held beliefs that a treatment will be effective (Berna et al., 2017; Vase et al., 2005). Patients’ expectations in augmenting placebo effects are perhaps most clearly observed in so-called “open-hidden” experimental investigations (Amanzio et al., 2001; Tondorf et al., 2017). For example, in an experiment in a hospital setting, intravenous analgesics were administered to patients covertly (from another room) or openly (in full view of the patient). Evidence for the influence of patients’ expectations was inferred from the finding that those

in the covert condition required a 50% higher dosage of analgesics to obtain the same relief as did those in the open condition (Amanzio et al., 2001).

However, it is also commonly theorized that expectancies can be shaped by other pathways, such as learning processes, particularly conditioning (Amanzio and Benedetti, 1999; Benedetti et al., 2003b); verbal instructions including the provision of a treatment rationale (Locher et al., 2017; Tondorf et al., 2017); and perceptions about others. Classical—or “Pavlovian”—conditioning refers to the learned associations between a neutral stimulus (e.g. a bell) and a biologically potent stimulus (e.g. food) to elicit a response (e.g. salivation) (Büchel et al., 2014; Kirsch et al., 2004; Rescorla, 1988). Its role in modulating placebo effects has been successfully demonstrated in a number of studies (Carlino et al., 2015; Colloca et al., 2006, 2008; Voudouris et al., 1985); research on conditioned placebo effects indicates further that the effects of conditioning may be mediated by response expectancies (Kirsch et al., 2014; Montgomery and Kirsch, 1997).

Verbal instructions can also influence expectancies about treatments. Recently, an innovative experiment in placebo analgesia found that placebos administered with a plausible rationale elicited significantly higher levels of pain-relieving placebo effects than placebos given without an explanation (Locher et al., 2017). In this study participants were prescribed “open label” placebos: that is, they were informed that the sugar pill might work by harnessing placebo effects. Crucially, in order for the open label placebos to work, it was not sufficient merely to prescribe placebos, clinicians had to offer a basic scientific rationale for their effectiveness (Locher et al., 2017).

Perceptions of clinicians’ competence and empathy are also likely to affect treatment outcome expectancies (Howe et al., 2017, 2019). Drawing on well-established social psychology research that perceptions of warmth and competence constitute important dimensions of interpersonal interactions (Fiske et al., 2002, 2007), Howe, Crum and colleagues argue that patients’ expectations about clinical interventions always

arise in a social context (Howe et al., 2017, 2019; Zion and Crum, 2018): “*When forming impressions, humans readily and rapidly determine whether another person’s intentions are benevolent (judgments of warmth), and whether this person has the ability to enact those intentions (judgments of competence)*” (Howe et al., 2017). Connecting these findings with research in placebo studies, it is hypothesized that patients’ perceptions both about clinician competence and warmth are key factors in establishing treatment expectancies, and inducing placebo effects (Howe et al., 2017, 2019; Zion and Crum, 2018). Put another way, placebo effects may be enhanced by patients’ perceptions that “the clinician gets it”—the practitioner displays knowledge, skill, proficiency, and personal effectiveness in relation to understanding the patients’ symptoms and condition; and “the clinician gets me” through signals of support, compassion, and personalized engagement (Howe et al., 2019). Providing an understandable treatment rationale may enhance perceptions of clinician competence via displays of expertise; and/or clinician warmth, by demonstrating careful articulation of the treatment rationale, and/or by investing time in patient-centered care.

Correlatively, nocebo effects are often described as “negative placebo effects” and are believed to be mediated by patients’ negative expectations about an intervention or a prognosis, resulting in adverse health responses, including pain, and with larger numbers of side effects associated with prescribed treatments (Benedetti et al., 2007; Colloca and Benedetti, 2007). Research into nocebo effects is more limited and there is no evidence that perceptions of low competence or low empathy, or lack of treatment rationale elicit these negative effects (Howe et al., 2017). Studies do suggest that adverse health effects are triggered by patients’ *negative* expectations about interventions (Barsky et al., 2002; Benedetti et al., 2003a). In a study of beta blockers prescribed for cardiac disease and hypertension, informing patients that side effects might include erectile dysfunction led to twice as many men reporting this problem, compared to those not informed (Silvestri et al., 2003). Experimental studies focusing on pain indicate that nocebo

effects engage specific regions of the brain involved in pain processing (Koyama et al., 2005). Open-hidden paradigms have also been applied to placebo studies with illuminating results (Benedetti et al., 2003a; Colloca et al., 2004). For example, in a study of postoperative pain among patients who received morphine for 48 hours, participants were allocated to either an open or hidden interruption in analgesic administration (Colloca et al., 2004). Those in the open condition were advised that morphine had been stopped; among patients in the hidden condition, morphine administration was stopped surreptitiously. After 10 hours, more patients in the open group requested additional pain-killers, suggesting that the verbal disclosure influenced expectations and, as a consequence, experiences of pain.

Hypotheses: How open notes might generate placebo and nocebo effects

Connecting evidence from placebo studies with the relatively novel platform of open notes, we propose that fully transparent patient portals may generate both placebo and nocebo effects.

Specifically, we hypothesize that patients may experience placebo effects under the following circumstances:

Hypothesis 1: Clinical notes convey positive expectations about the success of the treatment, and/or the patient's progress/prognosis; and/or

Hypothesis 2: Patients perceive the clinical notes to convey a persuasive rationale for treatment(s); and/or

Hypothesis 3: Patients perceive clinicians to be competent (e.g. the notes demonstrate complete and accurate information about the patients' conditions, and proposed treatments), and warm (e.g. the notes provide a high level of personal support, encouragement, and empathy for the patient's circumstances).

In addition, we propose that patients may experience nocebo effects if:

Hypothesis 4: Clinical notes convey negative expectations about the success of the treatment, including potential negative side effects.

We also suggest that there may be interactive effects between some of these hypothesized pathways. For example, communicating positive treatment outcome expectations may also enhance perceptions of clinician competence, thereby further boosting placebo effects.

Preliminary evidence that sharing notes might generate placebo and nocebo effects

Preliminary evidence for response expectancies

As conveyed in *Hypothesis 1*, we propose that the tone and content of clinical notes may play a direct role in influencing response expectancies among patients. By communicating encouraging and optimistic messaging anticipating the effectiveness of a treatment plan, and/or the patient's progress, there is potential to modulate the size of placebo effects among some patients. Currently, qualitative studies suggest that at least some individuals may experience positive emotions after reading their notes, and that the content of the note prompted these responses. For example: for example: "I enjoyed seeing my progress documented;" "Writes excellent very nice and specific notes that make me feel good, that I'm making progress with myself and that she sees the changes in me" (O'Neill et al., 2019). However, it is not understood whether these positive responses can be attributed to placebo effects, facilitated by reading clinical notes. Nonetheless, as a result of open notes some participants do appear to experience positive expectancies: for example: "I feel less helpless and perhaps more hopeful" (Gerard et al., 2017).

Preliminary evidence for provision of a treatment rationale

Studies show that many patients misunderstand or misremember what is communicated to them

about their medications and treatment plans (McCarthy et al., 2012). Survey evidence suggests that many patients report better understanding of treatment rationales a result of open notes, and this may provide a novel pathway toward eliciting placebo effects (Locher et al., 2017; Tondorf et al., 2017). For example, evidence from primary care indicates that patient understanding is enhanced by reading clinical notes (Esch et al., 2016). In a large survey across three disparate US health centers in 2017, access to notes enhanced patients' grasp of the rationale behind treatments and recommendations: 73% (16,354/22,520) rated reading their notes as very important for taking care of their health, 70% (15,726/22,515) as important for feeling more in control of their care, and 66% (14,821/22,516) for remembering their care plan. Only 3% of patients (737/22,304) reported being very confused after reading their notes (Walker et al., 2019). In the same survey, of 19,411 patients who read their notes and reported being prescribed medications, 14% responded that reading their notes made them more likely to take medications as prescribed (DesRoches et al., 2019).

Qualitative findings also suggest that reading clinical notes may improve grasp of treatment rationale. For example: *"I like knowing what the results of my tests mean. The records [laboratory results] show the numbers but the notes provide the interpretation in regards to my personal health status,"* *"I appreciate the open exchange and the opportunity to correct any possible misunderstandings,"* *"It is an opportunity to be more knowledgeable about my condition and how I can manage it better"* (Gerard et al., 2017).

Around a quarter of health organizations that share open notes in the US currently invite patients to view mental health notes. Many patients accessing mental health notes also report enhanced understanding about their condition. At the end of a 20-month long pilot study, 98% ($n = 44$) participants at an outpatient psychiatric clinic expressed a desire to continue reading their mental health notes online (Peck et al., 2017). Patients reported better understanding of

their mental health (69%, $n = 31$), and remembering their care plan (69%, $n = 31$) (Peck et al., 2017). In a study involving patient access to their psychotherapy notes ($n = 85$), more than half of patients rated notes as "very important" or "extremely important" for feeling more in control of their care (O'Neill et al., 2019). Although empowerment is a multi-dimensional concept in which patient knowledge is only one aspect, qualitative findings suggest that reading notes can improve understanding of what goes on in appointments and therapy sessions (Cromer et al., 2017; O'Neill et al., 2019; Peck et al., 2017). For example: *"It was confirming. It helped me understand my situation"; "helps affirm what I am working on"* (O'Neill et al., 2019).

By allowing patients greater time to read and reflect on what their clinician communicated away from the pressures of the face to face visit (Blease et al., 2020b), open notes may offer important opportunities to enhance response expectancies, and thereby facilitate placebo effects, via the provision of treatment rationales within online documentation.

Preliminary evidence for perceptions of clinician competence and warmth

To enhance positive expectations via the competence/warmth pathway (Howe et al., 2019), we predict that depending on the tone and content of the documentation, reading clinical notes might enhance patients' perceptions of clinician competence ("the clinician gets "it") and/or perceptions of clinician warmth ("the clinician gets me") (Howe et al., 2019). Many patients describe feeling empowered by reading their notes and report enhanced satisfaction levels with clinicians. After a year-long US pilot study ($n = 4592$), 37% of patients reported feeling better about their physician after reading their notes, with 62% expressing no difference (Bell et al., 2017). Around half (54%) of the primary care physicians ($n = 99$) believed that patient satisfaction with them had increased as a result of open notes, with similar numbers (51%, $n = 61$) believing that patients trusted them more (Bell et al., 2017). It is not yet

known whether increased levels of patient satisfaction are mediated by perceptions of clinician competence.

Comparable findings have emerged from studies of patients with mental illness. Interviews among Veterans Health Administration mental health clinicians and nurses believe that access to notes can empower patients, shift power dynamics in clinical sessions, and facilitate patient-centered care. If used carefully, open notes may enhance the therapeutic relationship (Denneson et al., 2017). To date, small sample surveys of access to mental health notes suggest that some patients express greater confidence and trust in their clinician as a result of reading their clinical notes (Cromer et al., 2017; O'Neill et al., 2019; Peck et al., 2017). For example, in a study of access to therapists' notes, one third of participants (32%, $n = 21$) described trusting their provider more, with 60% ($n = 39$) reporting no change (O'Neill et al., 2019). Again, while we might infer a connection between patient trust in clinicians and perceptions of clinician competence, there is no direct evidence that the former is affected by the latter. However, free text commentary accompanying surveys suggests that perceptions of clinicians' skills and abilities might be augmented after reading notes. For example: "[Reading the note] gave me insight into the evaluation process my doctor used and gave me confidence in his abilities (Bell et al., 2017); "I see how much my doctor really makes an effort to list and address my concerns" (Gerard et al., 2017).

Many patients express feelings of validation ("being heard") and of practitioner empathy. For example: "I felt like someone cared. May seem quite simple but it was a nice human touch" (Gerard et al., 2017); "I always appreciate how well my therapist captures what I've said and how I'm feeling" (O'Neill et al., 2019). Some describe an enhanced therapeutic alliance with their clinician. For example: "[The note] helps me feel that my [doctor] and I are partners in promoting my health" (Gerard et al., 2017), "I felt that my therapist was really listening to me" (O'Neill et al., 2019). Although tentative, these findings provide promising indications that cues of both empathy and warmth may already be communicated by some clinicians via clinical notes.

Preliminary evidence for nocebo effects

Open notes may also be a platform that produces negative expectancies and nocebo effects. In the large 2017 patient survey, of the 19,411 respondents who read their notes and were prescribed medications, 45% reported that they were more aware of the possible side effects of their prescriptions as a result (DesRoches et al., 2019a). A proportion of these respondents may also have derived nocebo effects from reading this information. In qualitative studies, some psychotherapy patients report feeling more negative about their progress as a result of accessing their clinician's notes. For example: "*The notes seemed separate from having a social worker as an ally in personal growth. I felt disempowered*", "*The therapist only said supportive things to me but the note seemed judgmental in a negative way. After reading it, I felt badly, like she didn't like me as much as I thought*" (O'Neill et al., 2019).

Negative expectancies may also arise if patients experience incongruencies between what is expressed by clinicians in face-to-face clinical encounters, and what is communicated in their notes (Cromer et al., 2017; O'Neill et al., 2019). For example: "*I felt uncomfortable that she told me one thing yet I read something else in the note. I don't know that I would see her again due to this*" (O'Neill et al., 2019).

Furthermore, in light of findings that access to notes may enhance understanding about treatments, patients may also become more vulnerable to nocebo effects. In a pilot study of access to psychiatric notes, Peck and colleagues found that most patients surveyed reported better understanding the potential side effects of their medications (82%, $n = 37$) (Peck et al., 2017). Whether enhanced understanding of possible side effects translates to increased nocebo effects has not yet been explored.

Future directions

Survey research provides a useful starting point for mounting hypotheses about placebo and

nocebo effects related to open notes. However, most surveys of patients' experiences of open notes in primary care contexts have been restricted to a few medical centers in the US, limiting the generalizability. In addition, only a small number of surveys have investigated access to psychiatric and psychotherapy notes, and in at least one of these studies, clinicians decided which patients should have access (Peck et al., 2017). As with all survey research, results are based on self-report, and responses may have been biased by individuals who were more engaged with patient portals and/or those who had more negative or positive experiences as a result of reading their clinical notes. Several RCTs examining the effects of sharing access to electronic health records have yielded positive results, but these studies are hampered by small sample sizes, and methodological limitations including the supplementation of patient portal access with clinician interventions (Goldzweig et al., 2013; Simon et al., 2011). Only few studies have investigated symptoms or conditions that are particularly responsive to placebo effects (Jones et al., 1999; Tuil et al., 2007).

To connect open notes research to research in placebo studies researchers will need to systematically collect well-validated measures of placebo/nocebo related processes such as treatment-related expectations. We suggest that future research into the connections between placebo and nocebo effects, and access to clinical notes should therefore encompass a range of novel approaches. First, independent thematic coding, the use of linguistic analysis software (Kahn et al., 2007; Pennebaker et al., 2007), or natural language processing (Rahimian et al., 2019), may be used to assess the length of notes, and their syntactic and semantic structure. Results of these findings might then be compared to patients' evaluations of their notes. This could help to probe whether objective linguistic markers are predictive of patients' responses—including expectations about treatments, perceptions of clinician competence and empathy, and understanding of treatment rationale. Where possible, we suggest that validated measures should be used to assess patients'

expectations about treatments, their subjective understanding of treatment rationales, and perceptions of clinician competence, and empathy. Importantly, although there are a number of instruments for assessing patient satisfaction with clinician communication, specific measures should be employed whenever possible, such as measuring of perceptions of clinician empathy following access to clinical notes.

Other clinical trials could compare different types of standardized or enhanced forms of open notes, and test how these are perceived by patients. For example, in one condition, clinicians could undergo training in clinical note-writing aimed at optimizing factors relevant to placebo effects. Clinicians might undertake a web-based course on patient-centered communication practices in open notes, aimed at improving understandable, supportive, and empathic writing skills (Dobscha et al., 2019). Similar to a placebo effect study conducted by Kaptchuk et al, clinicians could be tasked with incorporating several, specific cues into their clinical note-writing—such as an encouraging comment, a personal detail about the patient, an indication that the clinician understands the patient's health concerns, and a clear rationale for treatment recommendations (Kaptchuk et al., 2008). A second condition could involve a neutral, or no training clinician group. At set time periods, appraisals of patients' outcomes for placebo-effect responsive conditions could then be performed. For example, assessment of patients' primary symptoms, as well as adherence to treatments, could be measured, along with subjective patient health reports. We acknowledge, however, that setting up such a trial could be challenging since clinician blinding may not be possible. In addition, training in clinical note-writing might augment the quality of interpersonal care patient visits thereby interfering with accurate measures of the influence of documentation on placebo effects.

Beyond research into documentation, the very offer, or refusal, by clinicians to provide patients with ready access to their clinical notes may influence placebo and nocebo effects. While this issue is moot in some countries such as Sweden and the US where access to open

notes is mandated by law, conceivably in other regions even the act of inviting patients to read notes may increase response expectancies, and thereby placebo effects by enhancing trust in clinicians, and perceptions of clinician competence and empathy. On the other hand, explicit refusal to provide access to clinical notes may diminish clinician trust (O'Neill et al., 2019), driving negative expectations about prognoses and treatments, leading to nocebo effects. Both theories—that positive *and* negative decisions about providing access to clinical notes also may directly incur health effects—deserve exploration (DesRoches et al., 2019; Wright et al., 2015).

Finally, research might also venture beyond open notes to include not just the documentation but the medium through which it is shared (Gruszka et al., 2019; Torous and Firth, 2016). Evidence shows that users of digital devices often experience affiliative feelings, with many experiencing anxiety if separated from their smartphone (Clayton et al., 2015; Sapacz et al., 2016), and scales have been developed that intend to measure the therapeutic alliance that is formed with an application or program, rather than a person (Berry et al., 2018). We are aware of only one experiment that found placebo analgesia can be induced via the use of online communication (Pontén et al., 2019). Perceived sophistication of mobile Health (“mHealth”), encompassing the novelty and/or the design of apps, may increase expectations about the effectiveness of these interventions. Associative learning between neutral features of apps, or patient portals, and positive user experiences may engender conditioned placebo effects. Correlatively, some patients may learn to associate negative responses—for example, anxiety associated with distrust of technology—leading to negative conditioned responses upon accessing clinical notes via portals (Lopez et al., 2019).

Conclusion

Inviting patients to read their notes can be interpreted as a potential treatment tool—one that must be utilized with care. The hypotheses

proposed in this paper now require experimental research to investigate whether clinical notes can augment placebo and nocebo effects. We conclude with a final word of caution. Although desirable to maximize therapeutic benefits of placebo effects by communicating positive expectations to patients via clinical notes, this aspiration raises ethical considerations about upholding honesty in the disclosure of clinical information (Blease, 2012, 2019). If the hypotheses in this paper are empirically supported, it will be important to train clinicians to write notes that balance transparency (Blease et al., 2020b) with communication techniques that maximize the benefits of placebo effects, and minimize the harms of nocebo effects (Alfano, 2015; Blease, 2015; Evers et al., 2018; Fava et al., 2017; Klein et al., 2016).

Acknowledgements

The authors thank Ted Kaptchuk for helpful discussions on this topic.

Author contributions

Conceived the paper: CB, CD, JW. Wrote first draft: CB. Contributed to revisions and rewriting: IK, TD, JW, JT, MP, CD, MH.

Declaration of conflicting interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: CB and MH were supported by Keane Scholar Awards. CD, TD, JW were supported by the Gordon and Betty Moore Foundation, and the Cambia Health Foundation.

Statement of ethics

No ethical approval was required for this study.

ORCID iD

Charlotte R Blease  <https://orcid.org/0000-0002-0205-1165>

References

- Alfano M (2015) Placebo effects and informed consent. *The American Journal of Bioethics* 15(10), 3–12.
- Amanzio M and Benedetti F (1999) Neuropharmacological dissection of placebo analgesia: Expectation-activated opioid systems versus conditioning-activated specific subsystems. *Journal of Neuroscience* 19(1): 484–494.
- Amanzio M, Pollo A, Maggi G, et al. (2001) Response variability to analgesics: A role for non-specific activation of endogenous opioids. *Pain* 90(3): 205–215.
- Barsky AJ, Saintfort R, Rogers MP, et al. (2002). Nonspecific medication side effects and the nocebo phenomenon. *JAMA* 287(5): 622–627.
- Bell SK, Mejilla R, Anselmo M, et al. (2017). When doctors share visit notes with patients: A study of patient and doctor perceptions of documentation errors, safety opportunities and the patient–doctor relationship. *BMJ Qual Saf* 26(4): 262–270.
- Benedetti F, Lanotte M, Lopiano L, et al. (2007) When words are painful: Unraveling the mechanisms of the nocebo effect. *Neuroscience* 147(2): 260–271.
- Benedetti F, Maggi G, Lopiano L, et al. (2003a) Open versus hidden medical treatments: The patient’s knowledge about a therapy affects the therapy outcome. *Prevention & Treatment* 6(1): 1a.
- Benedetti F, Pollo A, Lopiano L, et al. (2003b) Conscious expectation and unconscious conditioning in analgesic, motor, and hormonal placebo/nocebo responses. *Journal of Neuroscience* 23(10): 4315–4323.
- Berna C, Kirsch I, Zion SR, et al. (2017) Side effects can enhance treatment response through expectancy effects: An experimental analgesic randomized controlled trial. *Pain* 158(6): 1014.
- Berry K, Salter A, Morris R, et al. (2018). Assessing therapeutic alliance in the context of mHealth interventions for mental health problems: Development of the mobile Agnew relationship measure (mARM) questionnaire. *Journal of Medical Internet Research* 20(4): e90.
- Blease C (2012). The principle of parity: The ‘placebo effect’ and physician communication. *Journal of Medical Ethics* 38(4): 199–203.
- Blease C (2015) Authorized concealment and authorized deception: Well-intended secrets are likely to induce nocebo effects. *The American Journal of Bioethics* 15(10): 23–25.
- Blease C (2018) Consensus in Placebo Studies: Lessons from The Philosophy of Science. *Perspectives in Biology and Medicine* 61(3): 412–429.
- Blease C and Annoni M (2019) Overcoming disagreement: A roadmap for placebo studies. *Biology & Philosophy* 34(2): 18.
- Blease C, Fernandez L, Bell SK, et al. (2020a) Empowering patients and reducing inequities: Is there potential in sharing clinical notes? *BMJ Quality & Safety*. Epub ahead of print March 2020. DOI: 10.1136/bmjqs-2019-010490.
- Blease CR (2019) The role of placebos in family medicine: ‘Implications of evidence and ethics for general practitioners’. *Australian Journal of General Practice* 48(10): 700.
- Blease CR, O’Neill S, Walker J, et al. (2020b) Sharing notes with mental health patients: Balancing risks with respect. *The Lancet Psychiatry*. Epub ahead of print 11 February 2020. DOI: 10.1016/S2215-0366(20)30032-8.
- Büchel C, Geuter S, Sprenger C, et al. (2014) Placebo analgesia: A predictive coding perspective. *Neuron* 81(6): 1223–1239.
- Carlino E, Torta DM, Piedimonte A, et al. (2015) Role of explicit verbal information in conditioned analgesia. *European Journal of Pain* 19(4): 546–553.
- Clayton RB, Leshner G and Almond A (2015) The extended iSelf: the impact of iPhone separation on cognition, emotion, and physiology. *Journal of Computer-Mediated Communication* 20(2): 119–135.
- Colagiuri B, Schenk LA, Kessler MD, et al. (2015) The placebo effect: From concepts to genes. *Neuroscience* 307: 171–190.
- Colloca L and Benedetti F (2007). Nocebo hyperalgesia: How anxiety is turned into pain. *Current Opinion in Anesthesiology* 20(5): 435–439.
- Colloca L, Benedetti F and Pollo A (2006) Repeatability of autonomic responses to pain anticipation and pain stimulation. *European Journal of Pain* 10(7): 659–665.
- Colloca L, Lopiano L, Lanotte M, et al. (2004) Overt versus covert treatment for pain, anxiety, and Parkinson’s disease. *The Lancet Neurology* 3(11): 679–684.
- Colloca L, Tinazzi M, Recchia S, et al. (2008) Learning potentiates neurophysiological and behavioral placebo analgesic responses. *Pain* 139(2): 306–314.
- Cromer R, Denneson LM, Pisciotto M, et al. (2017) Trust in mental health clinicians among patients

- who access clinical notes online. *Psychiatric Services* 68(5): 520–523.
- Delbanco T, Walker J, Bell SK, et al. (2012) Inviting patients to read their doctors' notes: A quasi-experimental study and a look ahead. *Annals of Internal Medicine* 157(7): 461–470.
- Denneson LM, Cromer R, Williams HB, et al. (2017) A qualitative analysis of how online access to mental health notes is changing clinician perceptions of power and the therapeutic relationship. *Journal of Medical Internet Research* 19(6): e208.
- DesRoches C, Leveille S, Bell SK, et al. (2020) The views and experiences of clinicians sharing medical record notes with patients. *JAMA Network Open* 3(3): e201753.
- DesRoches CM, Bell SK, Dong Z, et al. (2019) Patients managing medications and reading their visit notes: a survey of OpenNotes participants. *Annals of Internal Medicine* 171(1): 69–71.
- Dobscha SK, Denneson LM, Jacobson LE, et al. (2016) VA mental health clinician experiences and attitudes toward OpenNotes. *General Hospital Psychiatry* 38: 89–93.
- Dobscha SK, Kenyon EA, Pisciotta MK, et al. (2019) Impacts of a web-based course on mental health clinicians' attitudes and communication behaviors related to use of OpenNotes. *Psychiatric Services* 70(6): 474–479.
- Esch T, Mejilla R, Anselmo M, et al. (2016) Engaging patients through open notes: An evaluation using mixed methods. *BMJ Open* 6(1): e010034.
- Essén A, Scandurra I, Gerrits R, et al. (2018) Patient access to electronic health records: Differences across ten countries. *Health Policy and Technology* 7(1): 44–56.
- Evers AW, Colloca L, Blease C, et al. (2018) Implications of placebo and nocebo effects for clinical practice: Expert consensus. *Psychotherapy and Psychosomatics* 87(4): 204–210.
- Fava GA, Guidi J, Rafanelli C, et al. (2017) The clinical inadequacy of the placebo model and the development of an alternative conceptual framework. *Psychotherapy and Psychosomatics* 86(6): 332–340.
- Fiske ST, Cuddy AJ and Glick P (2007) Universal dimensions of social cognition: Warmth and competence. *Trends in Cognitive Sciences* 11(2): 77–83.
- Fiske ST, Cuddy AJ, Glick P, et al. (2002) A model of (often mixed) stereotype content: Competence and warmth respectively follow from perceived status and competition. *Journal of Personality and Social Psychology* 82(6): 878.
- Gerard M, Fossa A, Folcarelli PH, et al. (2017) What patients value about reading visit notes: A qualitative inquiry of patient experiences with their health information. *Journal of Medical Internet Research* 19(7): e237.
- Goldzweig CL, Orshansky G, Paige NM, et al. (2013) Electronic patient portals: Evidence on health outcomes, satisfaction, efficiency, and attitudes: A systematic review. *Annals of Internal Medicine* 159(10): 677–687.
- Gruszka P, Burger C and Jensen MP (2019) Optimizing expectations via mobile apps: A new approach for examining and enhancing placebo effects. *Frontiers in Psychiatry* 10: 365.
- Häggglund M, DesRoches C, Petersen C, et al. (2019) Patients' access to health records. *BMJ Clinical Research* 367: i5725.
- Howe LC, Goyer JP and Crum AJ (2017) Harnessing the placebo effect: Exploring the influence of physician characteristics on placebo response. *Health Psychology* 36(11): 1074.
- Howe LC, Leibowitz KA and Crum AJ (2019) When your doctor "Gets It" and "Gets You": The critical role of competence and warmth in the patient-provider interaction. *Frontiers in Psychiatry* 10: 475.
- Jensen KB, Kaptchuk TJ, Kirsch I, et al. (2012) Nonconscious activation of placebo and nocebo pain responses. *Proceedings of the National Academy of Sciences of the United States of America* 109(39): 15959–15964.
- Jones R, Pearson J, McGregor S, et al. (1999). Randomised trial of personalised computer based information for cancer patients. *BMJ* 319(7219): 1241–1247.
- Kahn JH, Tobin RM, Massey AE, et al. (2007) Measuring emotional expression with the Linguistic Inquiry and Word Count. *The American Journal of Psychology* 263–286.
- Kaptchuk TJ, Kelley JM, Conboy LA, et al. (2008) Components of placebo effect: Randomised controlled trial in patients with irritable bowel syndrome. *BMJ* 336(7651): 999–1003.
- Kaptchuk TJ and Miller FG (2015) Placebo effects in medicine. *New England Journal of Medicine* 373(1): 8–9.

- Kirsch I (1985) Response expectancy as a determinant of experience and behavior. *American Psychologist* 40(11): 1189.
- Kirsch I (1997) Response expectancy theory and application: A decennial review. *Applied and Preventive Psychology* 6(2): 69–79.
- Kirsch I (2018) Response expectancy and the placebo effect. In: *International Review of neurobiology* (Vol. 138). Amsterdam: Elsevier, pp. 81–93.
- Kirsch I (2019) Placebo effect in the treatment of depression and anxiety. *Frontiers in Psychiatry* 10.
- Kirsch I, Kong J, Sadler P, et al. (2014) Expectancy and conditioning in placebo analgesia: Separate or connected processes? *Psychology of Consciousness: Theory, Research, and Practice* 1(1): 51.
- Kirsch I, Lynn SJ, Vigorito M, et al. (2004) The role of cognition in classical and operant conditioning. *Journal of Clinical Psychology* 60(4): 369–392.
- Klein JW, Jackson SL, Bell SK, et al. (2016) Your patient is now reading your note: Opportunities, problems, and prospects. *The American Journal of Medicine* 129(10): 1018–1021.
- Koyama T, McHaffie JG, Laurienti PJ, et al. (2005) The subjective experience of pain: Where expectations become reality. *Proceedings of the National Academy of Sciences of the United States of America* 102(36): 12950–12955.
- Locher C, Nascimento A F., Kirsch I, et al. (2017) Is the rationale more important than deception? A randomized controlled trial of open-label placebo analgesia. *Pain* 158(12): 2320–2328.
- Lopez A, Schwenk S, Schneck CD, et al. (2019) Technology-based mental health treatment and the impact on the therapeutic alliance. *Current Psychiatry Reports* 21(8): 76.
- McCarthy DM, Waite KR, Curtis LM, et al. (2012) What did the doctor say? Health literacy and recall of medical instructions. *Medical Care* 50(4): 277.
- Mehta S, Jamieson T and Ackery AD (2019) Helping clinicians and patients navigate electronic patient portals: Ethical and legal principles. *Canadian Medical Association Journal* 191(40): E1100–E1104.
- Miller Jr DP, Latulipe C, Melius KA, et al. (2016) Primary care providers' views of patient portals: Interview study of perceived benefits and consequences. *Journal of Medical Internet Research* 18(1): e8.
- Montgomery GH and Kirsch I (1997) Classical conditioning and the placebo effect. *Pain* 72(1–2): 107–113.
- O'Neill S, Chimowitz H, Leveille S, et al. (2019) Embracing the new age of transparency: Mental health patients reading their psychotherapy notes online. *Journal of Mental Health* 28(5): 527–535.
- Peck P, Torous J, Shanahan M, et al. (2017) Patient access to electronic psychiatric records: A pilot study. *Health Policy and Technology* 6(3): 309–315.
- Pennebaker JW, Booth RJ and Francis ME (2007) *LIWC2007: Linguistic Inquiry and Word Count*. Austin, TX: Liwc. Net.
- Petersson L and Erlingsdóttir G (2018a) Open notes in Swedish psychiatric care (part 1): Survey among psychiatric care professionals. *JMIR Mental Health* 5(1): e11.
- Petersson L and Erlingsdóttir G (2018b) Open notes in Swedish psychiatric care (part 2): Survey among psychiatric care professionals. *JMIR Mental Health* 5(2), e10521.
- Pontén M, Ljótsson B and Jensen K (2019) Shaping placebo analgesic responses on the Internet: A randomized experimental trial. *Pain Reports* 4(3): e698.
- Rahimian M, Warner JL, Jain SK, et al. (2019) Significant and distinctive n-Grams in oncology notes: A text-mining method to analyze the effect of OpenNotes on clinical documentation. *JCO Clinical Cancer Informatics* 3: 1–9.
- Rescorla RA (1988) Pavlovian conditioning: It's not what you think it is. *American Psychologist* 43(3): 151.
- Richards T (2020) Light amid the gloom. *The BMJ Opinion*, 12 March. Available at: <https://blogs.bmj.com/bmj/2020/03/12/tessa-richards-light-amid-the-gloom/>
- Sapacz M, Rockman G and Clark J (2016) Are we addicted to our cell phones? *Computers in Human Behavior* 57: 153–159.
- Silvestri A, Galetta P, Cerquetani E, et al. (2003) Report of erectile dysfunction after therapy with beta-blockers is related to patient knowledge of side effects and is reversed by placebo. *European Heart Journal* 24(21): 1928–1932.
- Simon GE, Ralston JD, Savarino J, et al. (2011) Randomized trial of depression follow-up care by online messaging. *Journal of General Internal Medicine* 26(7): 698–704.

- Sugarman MA, Loree AM, Baltes BB, et al. (2014) The efficacy of paroxetine and placebo in treating anxiety and depression: A meta-analysis of change on the Hamilton Rating Scales. *PLoS One* 9(8): e106337.
- Tondorf T, Kaufmann L-K, Degel A, et al. (2017) Employing open/hidden administration in psychotherapy research: A randomized-controlled trial of expressive writing. *PLoS One* 12(11): e0187400.
- Torous J and Firth J (2016) The digital placebo effect: Mobile mental health meets clinical psychiatry. *The Lancet Psychiatry* 3(2): 100–102.
- Tuil WS, Verhaak CM, Braat DD, et al. (2007) Empowering patients undergoing in vitro fertilization by providing Internet access to medical data. *Fertility and Sterility* 88(2): 361–368.
- Vase L, Robinson ME, Verne GN, et al. (2005) Increased placebo analgesia over time in irritable bowel syndrome (IBS) patients is associated with desire and expectation but not endogenous opioid mechanisms. *Pain* 115(3): 338–347.
- Voudouris NJ, Peck CL and Coleman G (1985) Conditioned placebo responses. *Journal of Personality and Social Psychology* 48(1): 47.
- Walker J, Leveille S, Bell S, et al. (2019) OpenNotes after 7 years: Patient experiences with ongoing access to their clinicians' outpatient visit notes. *Journal of Medical Internet Research* 21(5): e13876.
- Weiss RD, O'malley SS, Hosking JD, et al. (2008) Do patients with alcohol dependence respond to placebo? Results from the COMBINE Study. *Journal of Studies on Alcohol and Drugs* 69(6): 878–884.
- Wolters F, Peerdeman KJ and Evers AW (2019) Placebo and nocebo effects across symptoms: From pain to fatigue, dyspnea, nausea and itch. *Frontiers in Psychiatry* 10: 470.
- Wright E, Darer J, Tang X, et al. (2015) Sharing physician notes through an electronic portal is associated with improved medication adherence: Quasi-experimental study. *Journal of Medical Internet Research* 17(10): e226.
- Zion SR and Crum AJ (2018) Mindsets matter: A new framework for harnessing the placebo effect in modern medicine. *International Review of Neurobiology* 138: 137–160.